

Post-Surgery Instructions EYELID SURGERY (Blepharoplasty)

Patient Name:	Date:
	Surgery Date:

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

NORMAL POST-OPERATIVE SYMPTOMS

Typical symptoms of eyelid surgery and signs to watch for include:

- Tightness in eyelid region and difficulty closing your eyes: Blurry vision, dry eye, burning, watery or itchy eyes.
 Bruising and swelling in the eyelid region. These are normal experiences as the skin, tissues and sensory nerves heal.
 Pain medication will help you cope with any discomfort. Consistent sharp pain should be reported to our office immediately
- **Asymmetry, the eyes look different, or heal differently.** The eyes may look or feel quite different from one another in the days following surgery. This is normal; no two eyes in nature or following surgery are perfectly symmetrical.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen eyelid or excessive bruising or fluid retention that is localized to one region.

To alleviate any discomfort, and to reduce swelling, you may apply cool (not cold) compresses to your eyes. Do not apply ice or anything frozen directly on the skin. Do not apply compresses to your cheeks. Soak soft plain white washcloths or gauze squares in ice water and wring out well. Apply directly to the eyelids, but do not apply any pressure. Apply cool compresses, for no longer than 20-minute intervals.

DAY OF SURGERY INSTRUCTIONS

your first days home from the hospital.	be clear to the adult who will monitor your
st is important in the early stages of healing, ear your own strength. Spend 10 minutes every	
ill be more comfortable for you, and can reduc	e swelling. Always keep your head elevated.
es: Do not wear contact lenses or eyeglasse atdoors or in any bright light.	es. Avoid bright light. Wear dark lens, large
following surgery. Stick to non-carbonated, no I water, milk and yogurt drinks. You must cons d for the first 24 hours. rescribed: If you have a pain pump, follow the d other medications you must take include:	ume at least 8 ounces of fluid every 2 hours.
mg	x per day
mg	x per day
ncisions will seep fluid and some blood for a she for cleansing incisions. Avoid alcohol and h	• •
i 11 1	Il be more comfortable for you, and can reduce see: Do not wear contact lenses or eyeglassed toors or in any bright light. following surgery. Stick to non-carbonated, not water, milk and yogurt drinks. You must consider for the first 24 hours. rescribed: If you have a pain pump, follow the dother medications you must take include: mg mg mg mcisions will seep fluid and some blood for a shape of the strength of the properties of the strength

TWO (2) TO SEVEN (7) DAYS FOLLOWING SURGERY

During this time you will progress with each day that p	asses. Ease into your daily activities.	You will receive clearance to begin driving
or return to work at your post-operative visit, or within	days.	
Your post-operative visit is scheduled for:		
	•	

- Continue to cleanse wounds as directed; you may shower. Take a warm, not hot shower. Do not rub your incisions.
- **Apply ointment and skincare as directed**. Do not use any glycolic, retinoid or other potentially irritating skincare products near your eyes or on your face.
- Take antibiotic medications and supplements as directed. Take pain medication only as needed. You may wish to switch from prescription pain medication to acetaminophen or ibuprofen.
- Continue to wear dark lens, large framed sunglasses whenever you are outdoors. You may begin wearing reading
 glasses as soon as it is comfortable for you. Do not wear soft contact lenses until you receive clearance to do so. Hard
 contact lenses may be difficult to remove. Do not attempt to wear these.
- Continue to keep your head elevated, including when sleeping.
- Do not resume any exercise other than regular walking. Walking is essential every day to prevent the formation of blood clots.
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

ONE (1) TO FOUR (4) WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- Continue your wound care as directed. Do not wear any makeup until all stitches are removed AND until your incisions no longer have any crusting or scabbing.
- Refrain from direct sun exposure. Continue to wear your sunglasses. If you are outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure. Your eyelids and face are highly susceptible to sunburn or the formation of irregular, darkened pigmentation.
- **Do not smoke.** While incisions may have healed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- Refrain from any strenuous exercise and from bending or lifting.
- You may begin sleeping in a modified reclining position. However do not sleep lying flat or on your stomach. If you are a side sleeper, two pillows under your head and a soft pillow under your mid-back and shoulders may offer more comfort.

Follow-up as directed. Your second post-operative visit is scheduled for:	

SIX (6) WEEKS FOLLOWING SURGERY

Healing will progress swelling and bruising continue to diminish.

- You may ease into your regular fitness routine. However uses of protective eyewear when outdoors and when swimming are essential.
- Discomfort or tightness and tingling in your eyelids will resolve.
- **No need to resume smoking.** You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

YOUR FIRST YEAR

- Continue good skincare and sun protection, healthy nutrition and fitness.
- Schedule any complementary procedures, as recommended. Botulinum injections or other treatments may be recommended to enhance your results, and to help your results to be long-lasting.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our office. Early
 intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after
 surgery.
- A one-year post surgery follow-up is recommended. However you may call our office at any time with your concerns or for needed follow-up.

	cial appearance may change too. You may wish to undergo revision surgery t life. Contact our office with any of your questions or concerns, at any time.
	ctions. I understand that following these instructions is solely my bility to ask my doctor and his or her staff any questions I have related d healing.
Patient Signature:	Date:
Printed Name of Patient:	
Signature of Practice Representative and Witness:	