

Post-Surgery Instructions AUTOLOGOUS FAT INJECTIONS

Patient Name:	Date:
	Surgery Date:

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

NORMAL POST-OPERATIVE SYMPTOMS

Whether this is your first procedure or a subsequent autologous fat injection treatment, follow these instructions precisely:

- If your injections are facial, do not bend over or lay flat following treatment. This may increase any swelling you may experience. Sleep with your head elevated for the first few days following treatment.
- If your lips have been injected, avoid pursing your lips for the first few days after injection. This means sipping from a cup rather than a straw, and no pursing a cigarette.

The following are normal experiences that can occur following treatment:

- Tenderness or bruising at the injection site may last a week or more.
- Redness and swelling are likely and may take 2 to 4 weeks or more to resolve.
- Asymmetry is possible: Opposite sides of your face, and different facial regions may react differently including swelling and bruising. This is normal. If asymmetry is severe, is accompanied by acute localized pain, or is not corrected within 7-10 days following your injection, contact our office immediately.
- If you develop any nodules, lumps or uneven appearance in the skin, please call our office immediately.

* IF THIS IS YOUR FIRST TREATMENT

You may also experience mild tenderness, swelling, bruising or discomfort at the site where fat was extracted prior to your injections. Treat these conditions in the same manner you will treat the fat injection sites.

To alleviate any discomfort, and to reduce potential swelling you may gently, and without pressure, apply cool, not cold compresses to the treatment site. Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses for no longer than 20-minute intervals.

DAY OF THE TREATMENT INSTRUCTIONS

You will only be released to the care of a re health and support you, around the clock in	sponsible adult. All of these instructions must be your first days home from the hospital.	oe clear to the adult who will monitor your			
 Rest, but not bed rest: While rest is important in the early stages of healing, equally important is that you are ambulated meaning that you are walking under your own strength. Spend 10 minutes every 2 hours engaged in light walking indoors you recover. Recline, do not lie down. This will be more comfortable for you, and can reduce swelling. Always keep your head elevated Do not bend forward or over. 					
Take all medication, exactly as p	orescribed: Oral pain medication, antibiotics ar	nd other medications you must take include:			
Antibiotic	mg	x per day			
Pain Medication	mg	x per day			
Other / Supplements					
cotton swab or gauze soaked in wa	sites clean. Injection sites may seep fluid and arm water is appropriate for cleansing. tly impair your safety prior to surgery and your a	•			
Relax. Do not engage in any stres you.	sful activities. Do not lift, push, or pull anything	. Take care of no one, and let others tend to			

TWO (2) TO SEVEN (7) DAYS FOLLOWING TREATMENT

During this time you will progress with each day that p	asses. Ease into your daily activities.	You will receive clearance to begin driving
or return to work at your post-operative visit, or within	days.	
Your post operative visit is esheduled for		
Your post-operative visit is scheduled for <mark>:</mark>	<u> </u>	

- Cleanse your skin as directed. You may shower. Take a warm, not hot shower. If you choose to use a hair dryer, use only the coolest setting and do not allow the compressed air to blow directly onto your face or other treated areas.
- Apply ointment and skincare as directed. Do not use any glycolic, retinoid or other potentially irritating skincare products on your face until you receive clearance to do so.
- Take antibiotic medications and supplements as directed. Take pain medication only as needed. You may wish to switch from prescription pain medication to acetaminophen or ibuprofen.
- Protect treated areas from sun exposure. Wear dark lens, large framed sunglasses and a wide-brimmed hat whenever you are outdoors. Wear an SPF 30 daily on both the treated area and the donor site.
- Continue to keep your head elevated, including when sleeping.
- Do not resume any exercise other than regular walking. Walking is essential every day to prevent the formation of blood
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

ONE (1) TO FOUR (4) WEEKS FOLLOWING TREATMENT

As you resume your normal daily activities, you must continue proper care and healing.

- Refrain from direct sun exposure. Continue to wear your sunglasses and a wide-brimmed hat. If you are outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure. The injected region and donor site are both highly susceptible to sunburn or the formation of irregular, darkened pigmentation.
- **Do not smoke** Smoking deprives your body of necessary oxygen that can result in poor healing
- a

•	be not smoke: officially deprives your body of necessary oxygen that can result in poor ficaling.			
•	You may ease into your regular fitness routine when you feel ready. However protective eyewear and a hat are necessary when outdoors.			
 You may begin sleeping in a modified reclining position. However do not sleep lying flat or on your stomach. If you a side sleeper, two pillows under your head and a soft pillow under your mid-back and shoulders may offer more comfort. 				
Follow	up as directed. Your second post-operative visit is scheduled for:			

SIX (6) WEEKS FOLLOWING SURGERY

Healing will progress; swelling continues to diminish.

- Follow-up treatment varies. Autologous fat is considered a permanent filler. However, some of the initial volume of
 fat injected will reabsorb and you may desire additional injections to achieve your results. Your results may be
 achieved following one treatment session. You may require more touch-up treatments or a subsequent injection
 session to achieve your final results. For this reason, some of your fat may be frozen and stored to use in
 subsequent treatments.
- **No need to resume smoking.** You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

YOUR FIRST YEAR

- Continue good skincare and sun protection, healthy nutrition and fitness.
- Schedule any complementary procedures, as recommended. Botulinum injections or specific skincare treatments may be recommended to enhance your results, and to help your results to be long-lasting.
- A one-year post treatment follow-up is recommended. However, you may call our office at any time with your concerns or for needed follow-up.

Your appearance will change with age. Your facial appearance may of procedures at a later date to help maintain your appearance throughout at any time.	
I have read and understand all of the above instructions. I understand that it is also my responsibility to ask to these instructions or about my procedure, health and healing.	•
Patient Signature:	Date:
Printed Name of Patient:	
Signature of Practice Representative and Witness:	