

## **Post-Surgery Instructions**

# NIPPLE & AREOLA RECONSTRUCTION

Patient Name:	Date:
	Surgery Date:

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

#### NORMAL POST-OPERATIVE SYMPTOMS

Typical symptoms and signs to watch for following nipple and areola reconstruction include the following:

- Tingling, burning, or intermittent shooting pain: These are normal experiences as the skin, tissues and sensory nerves
  heal. Pain medication and muscle relaxants will help you cope with any discomfort. Consistent sharp pain should be
  reported to our office immediately. You may also feel minor discomfort at the graft donor site of your nipple graft.
- Hypersensitivity of nipples or lack of sensitivity: This is normal and will gradually resolve over time. If this becomes
  painful or excessive notify our office immediately.
- Shiny skin or any itchy feeling: Swelling can cause the breasts' skin to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling of the breasts or at any nipple graft donor site. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. If the skin becomes red and hot to the touch, contact our office immediately.
- Asymmetry, the breasts look different, or heal differently: Breasts may look or feel quite different from one another in the days following surgery. This is normal; no two breasts are perfectly symmetrical in nature or following surgery.

#### CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen breast or bruising that is localized to one breast or region of the chest.

To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the breasts or any nipple graft site. Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses, for no longer than 20-minute intervals.

Do not apply heat.

## DAY OF SURGERY INSTRUCTIONS

you.

Whether you are releas	ed after surgery or after ar	n overnight stay in a	recovery center	or hospital, you	u will only be rele	eased to the care
of a responsible adult.	All of these instructions n	nust be clear to the	adult who will m	nonitor your he	alth and support	you, around the
clock in the first 24 hour	rs following surgery.					

	st is important in the early stages of healing, e r your own strength. Spend 10 minutes every	
Recline, do not lie down. This wi	Il be more comfortable for you, and can reduc	e swelling. Always keep your head elevated
	following surgery. Stick to non-carbonated, no water, milk and yogurt drinks. You must cons d for the first 24 hours.	•
Take all medication, exactly as p	rescribed: If you have a pain pump, follow the	e instructions specifically for your pain pump
· · · · · · · · · · · · · · · · · · ·	d other medications you must take include:	
Antibiotic	mg	x per day
Pain Medication	mg	x per day
Muscle Relaxant		
Other / Supplements		
a short time after surgery. Keep drecleansing incisions. Avoid alcohol bacterial ointment over the steri-stri  Wear your compression or elastic wraps only to cleanse your incision  Do not smoke. Smoking can great not smoke.	c wraps around the clock. Follow the instruct or to empty any drains.  tly impair your safety prior to surgery and you	preferably Dove or Ivory) is appropriate for steri-strips over your stitches. Apply antions specifically removing any compression ar ability to heal following surgery. You mus
Relax. Do not engage in any stress	ful activities. Do not lift, push, or pull anything.	Take care of no one, and let others tend to

### TWO (2) TO SEVEN (7) DAYS FOLLOWING SURGERY

During this time you will progress with each day that p	asses. Ease into your daily activities.	You will receive clearance to begin driving
or return to work at your post-operative visit, or within	days.	
Your post-operative visit is scheduled for:		

- Continue to cleanse wounds as directed; you may shower. Take a warm, not hot shower. Do not take a bath. Limit your shower to 10 minutes. Do not remove any steri-strips. Do not rub your incisions. Apply a fragrance free moisturizer to breast and surrounding skin, however not on your incisions.
  - **Take antibiotic medications and supplements as directed.** Take pain medication and muscle relaxants only as needed. You may wish to switch from prescription pain medication to acetaminophen or ibuprofen.
  - Continue to wear your bra around the clock.
  - Do not resume any exercise other than regular walking. Walking is essential every day to prevent the formation of blood clots.
  - Maintain a healthy diet. Do not smoke. Do not consume alcohol.

## ONE (1) TO FOUR (4) WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- **Discomfort or tightness and tingling will resolve.** Any lingering nipple sensitivity or lack of sensation should begin to greatly improve.
- Continue wound care as directed.
- Ease into your fitness routine. Avoid aerobic exercise that may cause a lot of bounce.
- **Do not smoke.** While incisions may have sealed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars. For your long-term health, there is no need to resume smoking.
- Continue to wear a proper support bra. You may sleep without a bra; however a camisole with built-in shelf support can be comfortable and provides added support as you continue to heal.
- You may begin sleeping flat. However do not sleep on your stomach. If you are a side sleeper, a soft pillow under your mid-back and shoulders may offer more comfort and support than a single pillow under your head.
- Practice good sun protection. Do not expose your breasts to direct sunlight. If you are outdoors, apply at least an SPF 30 to the chest area at least 30 minutes prior to sun exposure. Your chest region and breast skin are highly susceptible to sunburn or the formation or irregular, darkened pigmentation.

Follow-up as directed. Your second post-operative visit is scheduled for	

Tattooing to reconstruct your areola can be scheduled after your incisions have healed, or as directed by Dr. Antonetti.

## YOUR FIRST YEAR

- Practice monthly breast self-exam.
- Continue healthy nutrition, fitness, and sun protection.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our office. Early
  intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after
  surgery.
- A one-year post surgery follow-up is recommended. However, you may call our office at any time with your concerns or for needed follow-up.

Your body will change with age. The appearance of your breasts value date to help maintain your appearance throughout life. Contact	
I have read and understand all of the above instructions. I responsibility. I understand that it is also my responsibility to a to these instructions or about my procedure, health and healing	sk my doctor and his or her staff any questions I have related
Patient Signature:	Date:
Printed Name of Patient:	
Signature of Practice Representative and Witness:	